

Spectroscopic study of the sgB[e]-high mass X-ray binary CI Cam

A. Aret¹, M.L. Arias², A. Torres², L. Cidale², Tõnis Eenmäe¹

¹ *Tartu Observatory, University of Tartu, 61602 Tõravere, Tartumaa, Estonia*

² *Facultad de Ciencias Astronómicas y Geofísicas, UNLP and IALP (CONICET-UNLP), La Plata, Argentina*

³ *Astronomický ústav, Akademie věd České Republiky, Ondřejov, Czech Republic*

Contact / editor.baaa@gmail.com

Resumen / En este trabajo presentamos espectros ópticos e infrarrojos de la estrella binaria de rayos X CI Cam, compuesta por una estrella B[e] y un objeto compacto. Se observa variabilidad en la línea H α y en las líneas de He I y Fe II, que podría estar relacionada con el movimiento orbital y con una tercera componente del sistema.

Abstract / We present optical and near infrared spectra of the X-ray binary CI Cam. This system is formed by a compact object and a B[e] star. We report line profile variability in H α , He I and Fe II lines, that could be related with the orbital movement and the presence of a third component.

Keywords / binary stars, early-type emission stars, circumstellar matter

1. Introduction

CI Cameleopardalis is an eccentric binary system consisting of a B[e] star and a compact object, probably a white dwarf (e.g., Ishida, Morio & Ueda, 2004). The system has been identified as the optical counterpart of X-ray transient XTE J0421+560. Barsukova et al. (2006) obtained an orbital period of 19.41 days and an eccentricity of 0.62. First Gaia data release gave a parallax to CI Cam $\rho=0.71 \pm 0.24$ mas (Gaia Collaboration et al., 2016). Considering the observed magnitude $m_V=11.77$ and taking into account an extinction $E(B-V)=0.85$ (Robinson, Ivans & Welsh, 2002) we obtained an absolute magnitude $M_V=-1.69$ (+0.63 -0.90), which is in agreement with the spectral type B4 III-V[e] suggested by Barsukova et al. (2006).

2. CI Cam activity

CI Cam was the first galactic B[e] supergiant star detected during an X-ray outburst in 1998, revealing its binary nature. Few days after the outburst, the UB-VRI photometry started to decline and the IR spectra displayed H, He and Fe emission lines, as well as warm CO molecular emission bands in the 2.3-3.5 μm region (Clark et al., 1999, 2000). The molecular emission was proposed to arise due to collisional excitation from regions shielded from the stellar radiation, which requires high densities. Interferometric observations suggested that CI Cam developed an equatorial disk wind with a dust-free high-temperature zone close to the star viewed almost pole-on (e.g. Thureau et al., 2009). Over a period of 7 years the star kept stable and slightly brighter than in its pre-outburst state. Many years after the prominent outburst, Gemini/GNIRS and LBT-Luci I

K-band spectra revealed the absence of Pfund lines and CO molecular bands (Liermann et al., 2014) suggesting the loss/dilution of a high-density circumstellar material and the return of CI Cam to a quiescent transition phase.

Motivated by reports of a new potential outburst in 2016 (Wijngaarden et al., 2016), we spectroscopically monitored this star during several months. In spite of the fact that there was no real outburst, our observations provided valuable material for a deep study of CI Cam in its quiescent phase.

3. Optical observations

Follow-up optical spectroscopic observations from October 2016 to March 2017 were performed at 1.5-m telescope of the Tartu Observatory, Estonia using a single slit spectrograph attached to the Cassegrain focus. Spectra in the wavelength range 6315-6720 \AA with spectral resolution $R \simeq 10000$ were obtained using a 1800 lines mm^{-1} grating. Spectra with resolution $R \simeq 4500$ covering spectral interval from 6230 \AA to 6970 \AA were obtained with a 1200 lines mm^{-1} grating.

All data were reduced and corrected for heliocentric velocity using standard IRAF* tasks. Systemic velocity -51 km/s (Aret et al., 2016) was applied to all spectra.

An echelle spectrum with resolution $R \simeq 11000$, taken on November 20 2017, was obtained from BeSS database (observed by Olivier Thizy and reduced using ISIS** V5.8.0).

*IRAF was distributed by the National Optical Astronomy Observatories, which are operated by the Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the National Science Foundation.

**<http://www.astrosurf.com/buil/isis-software.html>

CI Cam spectroscopic variability




Figure 1: CI Cam Fe II emission line profiles for different epochs. Fe II 6416 Å (top left) and 6432 Å (top right), Fe II 6456 Å (bottom left) and 6516 Å (bottom right).




Figure 3: GEMINI/GNIRS medium resolution near-infrared spectra of CI Cam taken in 2016




Figure 2: $H\alpha$ (top) and He I 6678 Å (bottom) lines in velocity scale.

We searched for evidences of an outburst in $H\alpha$, He I and Fe II emission lines. In Figs.1 and 2 we show line profiles with spectral resolution $R \simeq 10\,000$ at different epochs. Spectra taken within one night revealed no variability beyond noise level and were combined for better signal-to-noise ratio. From the figures we can observe that between the end of 2016 and the beginning of 2017 CI Cam displayed no signs of outburst, the observed variability remained at the same level.

For comparison, we also show spectral line profiles from an earlier observation, which was obtained at Perek 2-m telescope at Ondřejov Observatory on September 30, 2011 using a Coude spectrograph with resolution $R \simeq 12\,000$ (Šlechta & Škoda, 2002).

4. Near infrared observations

The infrared spectroscopic data consists of JHK-band observations taken on November 5 and 14 and December 15, 2016 with GEMINI/GNIRS (Gemini Near Infrared Spectrograph). The spectra were obtained using the XD mode with the 0.15/pix short camera, the 111 l/mm grating and a 0.3 arcsec wide slit, that gives a spectral resolution of $R \approx 6000$. The grating was centered at two different wavelengths 2.13 μm and 2.32 μm , allowing to cover a spectral range between 9400 to 24000 Å. Similarly to what is observed in the optical range, IR features do not vary in a significant way during November and December 2016, ruling out the occurrence of an outburst. In Fig.3 we show parts of the near infrared spectra indicating the main identified features. The JHK spectra displays emission in hydrogen Paschen and Brackett series lines, and He I and Fe II lines, similar to the ones observed in the spectrum shortly after the 1998 outburst

(Clark et al., 1999). However, CO molecular emission, and He II, Na I, and Pfund series emission lines, that were detected by Clark et al. (1999), are not present in our spectra.

5. Comments and future work

CI Cam is a unique star that has shown spectroscopic and photometric variability in different timescales. Barsukova et al. (2006) reported variability in brightness and in the radial velocities of He I and Fe II lines. The orbital motion of the binary B[e] star + compact object is undoubtedly one of the causes of this variability, but not the only one. Barsukova et al. (2006) reported orbital period of 19.4 days, however Bartlett et al. (2019) found no evidence of such a period from X-ray observations. Based on photometric observations, Goranskij et al. (2017) suggested pulsations of the B-type star during CI Cam quiescence phase. Moreover, Goranskij & Barsukova (2009) proposed the presence of a third companion in the system. According to our new optical and infrared observations, although the outburst is not observed, all lines displayed short term variability that deserves to be studied in detail and discussed in the context of different hypothesis (pulsations, binary motion or the presence of a third component). Consequently, CI Cam is a star in permanent active state that deserves to be more deeply studied.

Acknowledgements: This research has used the NASA Astrophysics Data System (ADS) and the SIMBAD data base, operated at CDS, Strasbourg, France. This work has made use of the BeSS database, operated at LESIA, Observatoire de Meudon, France: <http://basebe.obspm.fr>

Part of this project has received funding from the European Union's Framework Programme for Research and Innovation Horizon 2020 (2014–2020), under the Marie Skłodowska-Curie grant Agreement No. 823734.

AA and TE acknowledge financial support from the institutional research funding IUT40-1 of the Estonian Ministry of Education and Research.

References

- Aret A., Kraus M., Šlechta M., 2016, MNRAS, 456, 1424
- Barsukova E.A., et al., 2006, Astronomy Reports, 50, 664
- Bartlett E.S., Clark J.S., Negueruela I., 2019, A&A, 622, A93
- Clark J.S., et al., 1999, A&A, 348, 888
- Clark J.S., et al., 2000, A&A, 356, 50
- Gaia Collaboration, et al., 2016, A&A, 595, A2
- Goranskij V.P., Barsukova E.A., 2009, Astrophysical Bulletin, 64, 50
- Goranskij V.P., et al., 2017, A. Miroshnichenko, S. Zharikov, D. Korčáková, M. Wolf (Eds.), *The B[e] Phenomenon: Forty Years of Studies*, Astronomical Society of the Pacific Conference Series, vol. 508, 307
- Ishida M., Morio K., Ueda Y., 2004, ApJ, 601, 1088
- Liermann A., et al., 2014, MNRAS, 443, 947
- Robinson E.L., Ivans I.L., Welsh W.F., 2002, ApJ, 565, 1169
- Šlechta M., Škoda P., 2002, Publications of the Astronomical Institute of the Czechoslovak Academy of Sciences, 90, 61
- Thureau N.D., et al., 2009, MNRAS, 398, 1309
- Wijngaarden M.J.P., et al., 2016, The Astronomer's Telegram, 9634